Huron River Watershed Council


Winter Stonefly Search 2013 Data



Protecting the river since 1965




Capniidae: slender winter stonefly




Nemouridae: broadback stonefly



Perlidae: Common stonefly



Perlodidae: Patterned stonefly



Taeniopterygidae: broadback winter stonefly

"X" indicates this family was found in the 2013 Stonefly Search

Abundance rating was subjectively assigned by the volunteers at the site (none, rare, some, frequent, abundant)

Sites highlighted in gray were not sampled this year.

SITE LOCATION	SITE	2013 SAMPLES				ALL YEA	RS	
	#	Capniidae Nemouridae Perlidae Perlodidae Taeniop- Abundan terygidae Rating		First sampled	# Times sampled	Avg. Family Count	Min Family Count	Max Family Count
Arms Creek: Walsh Road	1	Capnids and Taeniops are normally found here consistently, although were not found this year. x Rare	Taeniops 1	1996	12	1.8	1	3
Bancroft-Noles Drain: Lebo Park	89	Stoneflies have never been found here in January samples; the insectis poor the rest of the year as well.	community	2009	2	0.0	0	0
Boyden Creek: Delhi Rd	2	Capnids are found here consistently; other families are occasionally for this year). x	ound (but not	1995	13	1.3	1	2
Boyden Creek: Golf Course	3	Capnids are always found here; Perlodids and Taeniops are found oc	casionally.	1996	13	1.5	1	3
Boyden Creek: Huron River Drive	4	Stoneflies have never been found here, although the bug community it the rest of the year.	s pretty good	1995	11	0.0	0	0
Chilson Creek: Brighton Road	45	Stoneflies have only been found here twice (2007, 2011). None	0	1998	10	0.1	0	1
Chilson Creek: Chilson Road	5	Perlodids and Taeniops are found here consistently. This year, Capni found, making this the best sample ever at this site. x x Some	ds were also	1997	14	1.8	1	3
Davis Creek: 11 Mile Road	81	Stoneflies have never been found here in January samples. None	0	2004	7	0.0	0	0

SITE LOCATION	SITE	2013 SAMPLES			ALL YEA	A R S	
	#	Capniidae Nemouridae Perlidae Perlodidae Taeniop- Abundance Total #	First	# Times	Avg. Family		_
				sampled	Count	Count	Count
Davis Creek: Doane Road	6	Taeniops are always found here; Capnids and Perlodids are found in about 50% of the samples.	1996	14	2.0	1	3
Davis Creek: Pontiac Trail	7	Stoneflies are almost always found here. Only twice have stoneflies not been found (2004 and 2012).	1996	14	1.1	0	3
Davis Creek: Silver Lake Rd	49	Taeniops are always found here; Perlids are found here in about 50% of samples. This site and Mann Creek are the sites that have Perlids in January. x x Frequent 2	1998	12	1.7	1	2
Fleming Creek: Botanical Gardens	9	Like all of the Fleming Creek sites, Capnids are consistently found here; also this is the only site of the 5 on Fleming Creek at which Taeniops are found.	1995	12	1.6	1	2
Fleming Creek: Galpin Rd	84	Like all of the Fleming Creek sites, Capnids are consistently found here. Perlodids were also found here for the first time in 2013. x x Frequent 2	2005	7	1.1	1	2
Fleming Creek: Geddes Road	11	Like all of the Fleming Creek sites, Capnids are consistently found here, and Perlodids are often found, though not this year. x Abundant 1	1995	14	1.1	1	2
Fleming Creek: Radrick Farms	12	Like all of the Fleming Creek sites, Capnids are consistently found here.	1997	9	1.1	1	2
Fleming Creek: Warren Road	13	Like all of the Fleming Creek sites, Capnids are consistently found here. x Rare 1	1996	14	1.0	1	1
Greenoak Creek: Rushton Road	8	Stoneflies have never been found here in January samples; the insect community is poor the rest of the year as well.	1998	11	0.0	0	0
Hay Creek: M-36	15	Capnids and Taeniops are found here consistently, Perlodids have been found here in recent years.	1998	11	2.2	1	3
Honey Creek: Darwin Rd	16	Stoneflies are found here consistently.	1998	12	1.8	1	2
Honey Creek: Jackson Road	18	Capnids are found here about in about 50% of the samples historically; but they haven't been seen since 2008.	2000	12	0.5	0	1

SITE LOCATION	SITE	2013 SAMPLES			ALL YEA	A R S	
	#	Capniidae Nemouridae Perlidae Perlodidae Taeniop- Abundance Total #	First	# Times	Avg. Family		-
			sampled	sampled	Count	Count	Count
Honey Creek: Pratt Road	19	Stoneflies haven't been found since 2003; prior to this Capnids were found consistently.	1995	12	0.6	0	2
Tioney Grook. Trail Road	10		1000	12	0.0	Ū	_
		Capnids are found here consistently and Taeniops are often found.	4005	10			
Honey Creek: Wagner Road	20	x x Some 2	1995	13	1.4	1	2
Horseshoe Creek: Barker Road	98	This site was sampled for the first time this year; no stoneflies were found.	2012	1	0.0	0	0
Tioredenied Grook: Barker Read			2012	,	0.0		Ů
Hereater Court Mariti Barri	0.4	Taeniops are found here consistently; Capnids are found about 50% of the time	4000	40	4.4		0
Horseshoe Creek: Merrill Road	21	(including this year)	1996	13	1.4	1	2
		X Capnids are found here about 50% of the time.					
Hummocky Lick: M-36	63	Capital and round from about 50% of the time.	2001	10	0.5	0	2
		Capnids and Taeniops have both been found here in every sample since sampling					
Huron Creek: Dexter-Pinckney	22	began in 1995; this is the first year Perlodids were found.	1995	11	2.1	2	3
Training Crooks Boxes Timelary			1000			_	ŭ
		Taeniops are always found here; Capnids are found occasionally.					_
Huron River: Bell Road	62		2001	10	1.4	1	2
		Perlodids are always found here; 2011 marked the first sample with Taeniops.					
Huron River: Commerce Road	47		1998	10	1.0	0	2
		Since 2002, Capnids and Taeniops are found here sporadically. Prior to 2002, no					
Huron River: Cross Street	24	winter stoneflies were found.	1997	11	0.9	0	2
		x Rare 1					
		Capnids and Taeniops are found here consistently. Also, Perlodids were found here this year for the first time!					
Huron River: Flat Rock	23		2001	9	1.9	1	3
		x x Abundant 3					
		Taeniops are always found here; Perlodids have been found in 3 samples (2008,					
Huron River: Island Park	61	2011, 2013). Capniids are usually found; this was the first year that they were not	2001	11	2.2	2	3
		x x Some 2					
Huron River: Proud Lake Rec	64	Stoneflies have only been found here once- in 2003.	2001	9	0.1	0	1
		Techiana are almost always found here					
Huron River: US-23 (Liv. Co)	51	Taeniops are almost always found here.	2000	11	0.7	0	1
		x Rare 1					
		This is the best site that HRWC monitors during the spring and fall, but winter stonefly abundances are only modest. Perlodids and Taeniops are found here					
Huron River: White Lake Road	25	consistently.	1995	12	1.8	1	2
		x x Some 2					
		A A COMO Z					

SITE LOCATION	SITE	2013 SAMPLES			ALL YEA	A R S	
	#	Capniidae Nemouridae Perlidae Perlodidae Taeniop- Abundance Total #	First	# Times	Avg. Family		
Huron River: Zeeb Road	26	terygidae Rating Families Capnids and Taeniops are found here consistently, but Capnids were not found this year. However, this is the first time that Perlodids were found.	2000	sampled 12	Count 1.9	Count 1	Count 3
Livermore Creek: Doyle Road	27	This site was sampled for the first time in 2012; no stoneflies were found.	2012	1	0.0	0	0
Malletts Creek: Chalmers Rd	27	Stoneflies have never been found here in January samples; the insect community is poor the rest of the year as well.	1996	12	0.0	0	0
Malletts Creek: I-94	28	Stoneflies have never been found here in January samples; the insect community is poor the rest of the year as well.	1995	11	0.0	0	0
Malletts Creek: Main Street	56	Stoneflies have never been found here in January samples; the insect community is poor the rest of the year as well.	2000	9	0.0	0	0
Malletts Creek: Scheffler Park	29	Stoneflies have never been found here in January samples; the insect community is poor the rest of the year as well.	1997	9	0.0	0	0
Mann: VanAmberg Road	30	This is the best stonefly site that we monitor. Four families have been found here for 7 years in a row starting in 2007. x x x x Abundant 4	1996	13	2.8	1	4
Mill Creek: Fletcher Road	31	Capnids and Taeniops are found here often; though occasionally no stoneflies are found at all. x Rare 2	1996	12	1.7	0	3
Mill Creek: Ivey Road	32	Stoneflies are found here consistently.	1995	13	1.3	0	2
Mill Creek: Jackson Road	33	Capnids are always found here; Taeniops are often found here. x Frequent 2	1997	11	1.7	1	2
Mill Creek: Klinger Road		Capnids and Taeniops are found here often; ocassionally no stoneflies are found at all.	2001	8	1.3	0	2
Mill Creek: Letts at M-52	34	Capnids and Taeniops are found here consistently.	1995	15	1.4	0	2

SITE LOCATION	SITE	2013 SAMPLES			ALL YEA	RS	
	#	Capniidae Nemouridae Perlidae Perlodidae Taeniop- Abundance Total #	First	# Times	Avg. Family		-
		terygidae Rating Families	sampled	sampled	Count	Count	Count
Mill Creek: Manchester Road	55	Taeniops are almost always found here.	2000	10	0.9	0	2
Mill Creek: Parker Road	96	This site was sampled for the first time in 2012; no stoneflies were found.	2012	1	0.0	0	0
Mill Creek: Shield Road	80	Capnids and Taeniops are always found here.	2003	9	1.9	1	2
Mill Creek: Warrior Park	79	Capnids and Taeniops are always found here. x Abundant 2	2003	10	1.9	1	2
Millers Creek: Glazier Way	35	Stoneflies have only been found here once- in 2002. None 0	1997	14	0.1	0	2
Millers Creek: Plymouth Road	72	Stoneflies have never been found here in January samples; the insect community is poor the rest of the year as well.	2003	9	0.0	0	0
Narrow Gauge Creek: Green Rd	75	Capnids and Nemourids are often found here. Nemourids are rare and only found at one other sites during the Stonefly Search. This site is unusual; it is a tiny trickle of water seeping out of the ground.	2003	7	1.1	0	2
Norton Creek: West Maple Road	65	Stoneflies have never been found here in January samples; the insect community is poor the rest of the year as well. None 0	2001	10	0.0	0	0
Pettibone Creek: Commerce Road	67	Stoneflies have never been found here in January samples; the insect community is poor the rest of the year as well.	2002	7	0.0	0	0
Pettibone Creek: Livingston Road		In 2012, we found a Nemourid, and it was the first time stoneflies had been found at this site. Nemourids are rare in the Huron. The two places Nemourids have been found in the past were both streams with plenty of groundwater input.	2002	7	0.1	0	1
Port Creek: Armstrong Road	60	Stoneflies have never been found here in January samples; the insect community is poor the rest of the year as well. None 0	2000	7	0.0	0	0
Portage Creek: Dexter-Townhall Rd	37	Taeniops are found here consistently; Capniids and Perlodids are found here occasionally. x x 2 Abundant	1998	14	1.9	1	3
Portage Creek: Rockwell Road	94	This was the first year that this sites was sampled; no stoneflies found. 0 None	2013	0	0.0	0	0
		2 11600					

SITE LOCATION	SITE	2013 SAMP	LES					ALL YEA	RS	
	#	Capniidae Nemouridae Perlidae Perlodidae	Taeniop- terygidae	Abundance Rating	Total #	First sampled	# Times	Avg. Family Count	Min Family Count	Max Family Count
		This was the first year that this sites was sampl			7 41111100	campica	campica	Count	Count	Count
Portage Creek: Stockbridge	91		,							
				0	None	2013	1	0.0	0	0
Portage Creek: Unadilla	58	Stoneflies have only been found here once- in 2	2000.			2000	9	0.1	0	1
i ortage oreek. Orladilla	30			0	None	2000	3	0.1	O	'
		This site has been sampled twice; no stoneflies	vet							
Portage Creek: Williamsville	92	This site has been sampled twice, he storiomes	you.			2011	2	0.0	0	0
South Ore Creek: Bauer Road	52	Taeniops are almost always found here.				1999	11	0.8	0	2
										_
South Ore Creek: Hamburg Road	40	Taeniops are always found here; Capnids are for found for the first time this year.	ound occasi	ionally. Perlod	id were	1995	15	1.3	1	2
South Ore Creek. Hamburg Koau	40	x	x	Some	2	1995	13	1.3	'	2
		Stoneflies have never been found in January sa								
South Ore Creek: Lake Ridge	50	only mediocre during the rest of the year.				2000	9	0.0	0	0
		Stanofling house only been found here once in 1	008							
Swift Run: Shetland Rd	41	Stoneflies have only been found here once- in 1	990.			1995	12	0.2	0	2
				None	0					
Traver Creek: Broadway	42	Capnids are found here about every other year.				1995	13	0.6	0	1
,		х		Rare	1					
		Capnids were found here in every sample from found for 2 samples (2009, 2011), but were found								
Traver Creek: Dhu Varren Rd	43	population still seems low.	ia again iiii	o your, aitriou	gii (ii)	1997	12	0.9	0	2
		х								
- 0 1		This is the first time that this site has been sam	pled.							
Traver Creek: Traver Road	101	x		Rare	1	2013	1	1.0	1	1
		Stoneflies have never been found here in Janua	ary samples							<u>_</u>
Walker Creek: 8 Mile Road	82					2004	6	0.0	0	0
		2011 was the first time this site was sampled, n	o stoneflies	were found						
Willow Run: Van Buren Park	90					2011	1	0.0	0	0

SITE LOCATION	SITE	2013 SAMPLES	S			ALL YEARS				
	#	l '	eniop- / /gidae	Abundance Rating	Total # Families	First sampled	# Times sampled	Avg. Family Count	Min Family Count	Max Family Count
Woodruff Creek: Buno Road	46	Stoneflies are only found here in about 1 of every 2 swe found 2 families for the first time!	samples.	. This year t	hough, 2	1995	11	0.4	0	2
Woodruff Creek: Maxfield Road	44	Stoneflies have been found here only twice since 19	997.	None	0	1997	12	0.2	0	1
Woods Creek: L Huron Metropark		Capnids, Perlodids, and Taeniops are typically found first year in 12 years that no stoneflies were found. S		lowever, this None	is the	1997	12	1.6	0	3
Woods Creek: Martinsville Road	87	Stoneflies have never been found here in January sa	samples.	None	0	2009	2	0.0	0	0
Woods Creek: Renton Road	88	2012 marked the first time that stoneflies were found	nd at this s	site!		2009	4	0.3	0	1

SITE LOCATION	SITE	2013 SAMPLES			ALL YEARS					
	#	Capniidae Nemouridae Perlidae Perlodidae	Taeniop-	Abundance	Total #	First	# Times	Avg. Family	Min Family	Max Family
			terygidae	Rating	Families	sampled	sampled	Count	Count	Count

 -		